



AMERICAN SOCIETY OF
PLASTIC SURGEONS®

Pre-Operative Patient Information



Please complete the following information prior to surgery.

Drug Reactions / Allergies / Latex Sensitivity

Some patients cannot take certain medications such as penicillin because of allergic reactions. Other patients experience reactions such as nausea / vomiting from narcotic pain medications (Codeine, Morphine, Demerol, Vicodin, Percocet, etc.). Please list below regarding any known drug allergies or reactions, or sensitivities.

Medication Name	Type of Drug Reaction / Allergy
1. _____	_____
2. _____	_____
3. _____	_____
4. _____	_____

_____ I do not have known drug allergies, drug reactions, or latex sensitivity.

Prescription Medications

Please list all prescription medications you currently take:

1. _____
2. _____
3. _____
4. _____

_____ I am not currently taking any prescription medications.

Non-Prescription Medications / Dietary Supplements / Vitamins / "Herbs" / Minerals

Many patients take non-prescription medications such as aspirin, anti-inflammatories (Advil, Motrin, Aleve) and other preparations that can be purchased without a prescription (dietary supplements, vitamins, "herbs", and minerals). Many of these can have profound effects on increased risk of bleeding during and after surgery or react with prescription medications. If you currently take items in this category, please list below. Please discontinue taking all non-prescription medications, dietary supplements, vitamins, herbs, and minerals for a minimum of 10 days before and after surgery.

1. _____
2. _____
3. _____

_____ I am not currently taking non-prescription medications, dietary supplements, vitamins, herbs, or minerals.

PRE-OPERATIVE PATIENT INFORMATION

Smoking, Second-Hand Smoke Exposure, Nicotine Products (Patch, Gum, Nasal Spray)-

Patients who are currently smoking, use tobacco products, or nicotine products (patch, gum, or nasal spray) are at a greater risk for significant surgical complications of skin dying and delayed healing. Individuals exposed to second-hand smoke are also at potential risk for similar complications attributable to nicotine exposure. Additionally, smoking may have a significant negative effect on anesthesia and recovery from anesthesia, with coughing and possibly increased bleeding. Individuals who are not exposed to tobacco smoke or nicotine-containing products have a significantly lower risk of this type of complication. Please indicate your current status regarding these items below:

_____ I am a non-smoker and do not use nicotine products. I understand the risk of second-hand smoke exposure causing surgical complications.

_____ I am a smoker or use tobacco / nicotine products. I understand the risk of surgical complications due to smoking or use of nicotine products.

_____ I have smoked and stopped approximately _____ ago. I understand I may still have the effects and therefore risks from smoking in my system, if not enough time has lapsed.

Date/Time: _____

Signature: _____